


5. Geometric Transformations and Projections

5.1 Translations and Rotations

a) Translation


$$\mathbf{P}_1 = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix}$$

$$\mathbf{P}_2 = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} + \begin{bmatrix} x_d \\ y_d \\ z_d \end{bmatrix} = \begin{bmatrix} x_1 + x_d \\ y_1 + y_d \\ z_1 + z_d \end{bmatrix} = \mathbf{P}_1 + \mathbf{d}$$




b) Scaling

$$\mathbf{P}_1 = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} \quad \mathbf{P}_2 = \begin{bmatrix} sx_1 \\ sy_1 \\ sz_1 \end{bmatrix} \quad \mathbf{P}_2 = s\mathbf{P}_1$$




c) Reflection (about x - 0 - y Plane)

$$\mathbf{P}_1 = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} \quad \mathbf{P}_2 = \begin{bmatrix} x_1 \\ y_1 \\ -z_1 \end{bmatrix} \quad \mathbf{P}_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$



d) Rotation about z Axis

$$\begin{aligned}\mathbf{P}_1 &= \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = \begin{bmatrix} r \cos f \\ r \sin f \\ z \end{bmatrix} \\ \mathbf{P}_2 &= \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} r \cos(f+q) \\ r \sin(f+q) \\ z \end{bmatrix} \\ &= \begin{bmatrix} r \cos f \cos q - r \sin f \sin q \\ r \cos f \sin q + r \sin f \cos q \\ z \end{bmatrix} \\ &= \begin{bmatrix} x_1 \cos q - y_1 \sin q \\ x_1 \sin q + y_1 \cos q \\ z_1 \end{bmatrix}\end{aligned}$$



5.2. Homogeneous Representation

The representation is introduced to express all geometric transformations in the form of matrix multiplication for the convenience of manipulation.

a) Translation

$$\mathbf{P}_2 = [\mathbf{D}] \mathbf{P}_1 \text{ or } \begin{bmatrix} x_2 \\ y_2 \\ z_2 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & T_x \\ 0 & 1 & 0 & T_y \\ 0 & 0 & 1 & T_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ y_1 \\ z_1 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + T_x \\ y_1 + T_y \\ z_1 + T_z \\ 1 \end{bmatrix}$$

b) Scaling

$$\mathbf{P}_2 = [\mathbf{S}] \mathbf{P}_1 \text{ and } [\mathbf{S}] = \begin{bmatrix} S_x & 0 & 0 & 0 \\ 0 & S_y & 0 & 0 \\ 0 & 0 & S_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

c) Reflection (Mirroring)

$$\mathbf{P}_2 = [\mathbf{M}] \mathbf{P}_1 \text{ and } [\mathbf{M}] = \begin{bmatrix} \pm 1 & 0 & 0 & 0 \\ 0 & \pm 1 & 0 & 0 \\ 0 & 0 & \pm 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


d) Rotation about the z Axis

$$\mathbf{P}_2 = [R_z] \mathbf{P}_1 \quad \text{and} \quad [R_z] = \begin{bmatrix} \cos\theta & -\sin\theta & 0 & 0 \\ \sin\theta & \cos\theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Similarly


$$[R_x] = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\theta & -\sin\theta & 0 \\ 0 & \sin\theta & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}; \quad [R_y] = \begin{bmatrix} \cos\theta & 0 & \sin\theta & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\theta & 0 & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Rotation about Z Axis - CCW by θ




$$\begin{aligned} \mathbf{P}_1 &= \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = \begin{bmatrix} r \cos f \\ r \sin f \\ z_1 \end{bmatrix} \\ \mathbf{P}_2 &= \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} r \cos(f+q) \\ r \sin(f+q) \\ z_1 \end{bmatrix} = \begin{bmatrix} r \cos f \cos q - r \sin f \sin q \\ r \cos f \sin q + r \sin f \cos q \\ z_1 \end{bmatrix} \\ &= \begin{bmatrix} x_1 \cos q - y_1 \sin q \\ x_1 \sin q + y_1 \cos q \\ z_1 \end{bmatrix} \\ \mathbf{P}_2 &= \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} \cos q & -\sin q & 0 \\ \sin q & \cos q & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = R_{[z]} \mathbf{P}_1 \end{aligned}$$

Rotation about X Axis - CCW by θ




$$\begin{aligned}
 \mathbf{P}_1 &= \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = \begin{bmatrix} x_1 \\ r \cos f \\ r \sin f \end{bmatrix} \\
 \mathbf{P}_2 &= \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} x_1 \\ r \cos(f+q) \\ r \sin(f+q) \end{bmatrix} = \begin{bmatrix} x_1 \\ r \cos f \cos q - r \sin f \sin q \\ r \cos f \sin q + r \sin f \cos q \end{bmatrix} \\
 &= \begin{bmatrix} x_1 \\ y_1 \cos q - z_1 \sin q \\ y_1 \sin q + z_1 \cos q \end{bmatrix} \\
 \mathbf{P}_2 &= \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos q & -\sin q \\ 0 & \sin q & \cos q \end{bmatrix} \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = R_{[x]} \mathbf{P}_1
 \end{aligned}$$

Rotation about Y Axis - CCW by θ



$$\begin{aligned}
 \mathbf{P}_1 &= \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = \begin{bmatrix} r \cos f \\ y_1 \\ r \sin f \end{bmatrix} \\
 \mathbf{P}_2 &= \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} r \cos(f - q) \\ y_1 \\ r \sin(f - q) \end{bmatrix} = \begin{bmatrix} r \cos f \cos q + r \sin f \sin q \\ y_1 \\ r \sin f \cos q + r \cos f \sin q \end{bmatrix} \\
 &= \begin{bmatrix} x_1 \cos q + z_1 \sin q \\ y_1 \\ -x_1 \sin q + z_1 \cos q \end{bmatrix} \\
 \mathbf{P}_2 &= \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} \cos q & 0 & \sin q \\ 0 & 1 & 0 \\ -\sin q & 0 & \cos q \end{bmatrix} \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = R_{[y]} \mathbf{P}_1
 \end{aligned}$$

Rotation about an Arbitrary Axis




Parametric Rep. of the Axis:


$$\begin{aligned}
 x &= Au + x_0 \\
 y &= Bu + y_0 \\
 z &= Cu + z_0 \quad 0 < u < 1 \\
 L &= \sqrt{A^2 + B^2 + C^2} u
 \end{aligned}$$

Step 1: Translate \mathbf{P}_0 to Origin

$$[D] = \begin{bmatrix} 1 & 0 & 0 & -x_0 \\ 0 & 1 & 0 & -y_0 \\ 0 & 0 & 1 & -z_0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$



Step 2: Rotate Vector \bar{L} about X Axis to get \bar{L} into the x - z plane



$$L = \sqrt{A^2 + B^2 + C^2}$$

$$V = \sqrt{B^2 + C^2}$$


$$\sin \theta_1 = \frac{B}{V}$$

$$\cos \theta_1 = \frac{C}{V}$$

$$[R_x] = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta_1 & -\sin \theta_1 & 0 \\ 0 & \sin \theta_1 & \cos \theta_1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{C}{V} & -\frac{B}{V} & 0 \\ 0 & \frac{B}{V} & \frac{C}{V} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Step 3: Rotate \bar{L} about the Y axis to get it in the z direction

Rotate a negative angle (CW)!




$$\sin \theta_2 = -\frac{A}{L}$$

$$\cos \theta_2 = \frac{V}{L}$$

$$[R_y] = \begin{bmatrix} \cos \theta_2 & 0 & \sin \theta_2 & 0 \\ 0 & 1 & 0 & 0 \\ -\sin \theta_2 & 0 & \cos \theta_2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{V}{L} & 0 & -\frac{A}{L} & 0 \\ 0 & 1 & 0 & 0 \\ \frac{A}{L} & 0 & \frac{V}{L} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Step 4: Rotate angle θ about axis \bar{L} .



$$[R_z] = \begin{bmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Step 5: Reverse the rotation about the Y axis

$$[R_y]^{-1} = \begin{bmatrix} \frac{V}{L} & 0 & \frac{A}{L} & 0 \\ 0 & 1 & 0 & 0 \\ -\frac{A}{L} & 0 & \frac{V}{L} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Inverse of Rotation:

Replace θ by $-\theta$

$\sin \theta$ by $-\sin \theta$

$\cos \theta$ by $-\cos \theta$


Step 6: Reverse rotation about the X axis

$$[R_x]^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{C}{V} & \frac{B}{V} & 0 \\ 0 & -\frac{B}{V} & \frac{C}{V} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Step 7: Reverse translation

$$[D]^{-1} = \begin{bmatrix} 1 & 0 & 0 & x_0 \\ 0 & 1 & 0 & y_0 \\ 0 & 0 & 1 & z_0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Overall Transformation



$$[T] = [D]^{-1} [R_x]^{-1} [R_y]^{-1} [R_z^q] [R_y] [R_x] [D]$$


$$\mathbf{P}_2 = [T]\mathbf{P}_1$$

Homework:


1. Give real X_0, Y_0, Z_0, A, B, C and theta values, find $[T]$.
2. A line connects the point A at $(1,0,0)$ to the point B at $(1,0,1)$. A second line extends from C at $(1,0,2)$ to D at $(1,1,2)$. Rotate line AB about line CD using vector-matrix methods. The rotation should be 90° counter-clockwise as seen from the +y axis.
3. A plane surface intersects the coordinate axes at three points $A=(5,0,0)$, $B=(0,5,0)$ and $C=(0,0,10)$. A given point P is on the plane. Find the matrix of geometric transformation that move the point P five units down on the plane to P' . (Line PP' is perpendicular to edge AB and is on plane ABC. $PP'=5$.)

5.3 Viewing Coordinate System


a) View Window



- Right hand coordinate system
- 2D coordinate system on the view plane.
- Some "old" graphics systems use a left-hand coordinate system. A different geometric transformation matrix must be used.




b) Orthographic View versus World Coordinate System



Isometric Projection

How to get it?




c) Perspective and Parallel Projections

We want to produce a 2D image (projection) of a 3D object.


- Perspective Projection
- Parallel Projection

a 3D object.

① Perspective Projection



② Parallel Projection




center of
projection, CP
at infinity

* Parallel lines appear as
parallel lines.

- Examples of Perspective Projection

5.4 Definition of a General Viewing Coordinate System

- a) What are involved?



- b) How to Set Up the Viewing Coordinate System (VCS)!

- i) Define the view reference point

$$\mathbf{P} = (P_x, P_y, P_z)^T$$

- ii) Define the line of the sight vector \bar{n} (normalized)

$$\bar{n} = (N_x, N_y, N_z)^T \text{ and } N_x^2 + N_y^2 + N_z^2 = 1$$

- iii) Define the "up" direction


$$\bar{V} = (V_x, V_y, V_z)^T \perp \bar{n}, \quad \bar{V} \cdot \bar{n} = 0$$

This also defines an orthogonal vector, \bar{u}

$$\bar{u} = \bar{V} \times \bar{n}$$

$(\bar{u}, \bar{V}, \bar{n})$ forms the viewing coordinates

iv) Define the View Window in $\bar{U} - \bar{V} - \bar{W}$ coordinates



c) Parallel Projection

First transform coordinates of objects into the UVn coordinates (VCS), then drop the n component. (n – depth)

Overlapping x - y - z and U - V - n

i) Translate O_v to O.

$$[D] = \begin{bmatrix} 1 & 0 & 0 & -0_{vx} \\ 0 & 1 & 0 & -0_{vy} \\ 0 & 0 & 1 & -0_{vz} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

ii) Align the \bar{n} axis with the Z axis.


$$A = N_x, \quad B = N_y, \quad C = N_z$$

$$L = \sqrt{N_x^2 + N_y^2 + N_z^2}$$

$$V = \sqrt{N_y^2 + N_z^2}$$

The procedure is identical to that given in 5.2.


• rotate θ_1 about x: $[R_x]$



• rotate θ_2 about y: $[R_y]$

iii) Rotate θ_3 about the Z axis to align \bar{U} with x and/or \bar{V} with y. At this point, \bar{V} is given by $(V'_x, V'_y, 0)^T$ where


$$\begin{pmatrix} V'_x \\ V'_y \\ 0 \\ 1 \end{pmatrix} = [R_y] [R_x] [D_{Ov, o}] \begin{pmatrix} V_x \\ V_y \\ V_z \\ 1 \end{pmatrix}$$



We need to rotate by an angle θ_3 about the Z axis

$$L = \sqrt{V'_x^2 + V'_y^2}, \quad \sin \theta_3 = \frac{V'_x}{L}, \quad \cos \theta_3 = \frac{V'_y}{L}$$

$$[R_z] = \begin{bmatrix} V'_{y/L} & -V'_{x/L} & 0 & 0 \\ V'_{x/L} & V'_{y/L} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$




iv) Drop the \mathbf{n} coordinate

$$[D_n] = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad \begin{pmatrix} u \\ v \\ 0 \\ 1 \end{pmatrix} = [D_n] \begin{pmatrix} u \\ v \\ n \\ 1 \end{pmatrix}$$

In summary, to project a view of an object on the UV plane, one needs to transform each point on the object by:

$$[T] = [D_n] [R_z] [R_y] [R_x] [Do_v, o]$$

$$\mathbf{P}' = \begin{pmatrix} u \\ v \\ 0 \\ 1 \end{pmatrix} = [T]\mathbf{P} = [T] \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$



Note: The inverse transforms are not needed! We don't want to go back to x - y - z coordinates.